



**Annual meeting of the  
Croatian Immunological Society  
2019**



**Rovinj  
11-12.10.2019**

Diamond Sponsor



## THE ROLE OF *MID1-PP2A* AXIS IN INFLAMMATION-INDUCED JOINT DAMAGE IN ANTIGEN-INDUCED ARTHRITIS

Nina Lukač<sup>1,2</sup>, Darja Flegar<sup>1,3</sup>, Tomislav Kelava<sup>1,3</sup>, Alan Šučur<sup>1,3</sup>, Katerina Zrinski Petrović<sup>1,2</sup>, Dino Šiš<sup>1,3</sup>, Maša Filipović<sup>1,3</sup>, Vedran Katavić<sup>1,2</sup>, Danka Grčević<sup>1,3</sup>, Nataša Kovačić<sup>1,2</sup>

1 Laboratory for Molecular Immunology, Croatian Institute for Brain Research, University of Zagreb, School of Medicine, Zagreb, Croatia

2 Department of Anatomy, University of Zagreb School of Medicine, Zagreb, Croatia

3 Department of Physiology and Immunology, University of Zagreb School of Medicine, Zagreb, Croatia

**INTRODUCTION:** Rheumatoid arthritis (RA) is a chronic autoimmune joint disease which often causes structural joint damage not reversible by currently available therapeutics, urging discovery of new molecular mediators for therapeutic targeting. In a murine model of RA, antigen-induced arthritis (AIA) we have previously shown that destructive AIA is characterized by accumulation of synovial myeloid cells which up-regulate *Mid1* gene. Due to the previously described involvement of *Mid1* in pathogenesis of immune-mediated diseases, the objective of this study was to evaluate the role of *Mid1* in AIA.

**MATERIALS AND METHODS:** AIA was induced in C57BL6 mice by immunization with methylated bovine serum albumin (mBSA) and subsequent intra-articular injection of mBSA. Expression of *Mid1* and pro-inflammatory cytokines was assessed by qRT-PCR. Mice were treated with metformin, which prevents *Mid1*-mediated ubiquitination of PP2A, and thus has an anti-inflammatory effect, at a daily dose of 1g/kg, to evaluate the effects of blocking the *Mid1*-mediated inflammation on arthritis development and subchondral bone resorption (assessed by  $\mu$ -CT).

**RESULTS:** *Mid1* was up-regulated in knee joints early after arthritis induction. Its expression positively correlated with severity of arthritis assessed by knee diameter measurement and levels of pro-inflammatory cytokines, IL-1, IL-6 and TNF. Although *Mid1* is positioned on the X-chromosome, its expression was up-regulated in arthritic joints of both male and female mice. Metformin treatment ameliorated the severity of arthritis and arthritis-induced subchondral bone resorption.

**CONCLUSION:** Inhibition of *Mid1*-mediated PP2A degradation by metformin might aid to therapeutic management of inflammatory arthritides.

**SUPPORT:** Croatian Science Foundation projects IP-2014-09-7406, DOK-2015-10-9897, IP-2018-01-2414.